

Contact Interactions of Free Anyons

S. VOROPAEV

VERNADSKY INSTITUTE, LAB.THEORETICAL AND MATHEMATICAL PHYSICS, RUSSIA
ACADEMY OF SCIENCE, KOSYGINA ST. - 19, MOSCOW

"Free" (non-scattered) Schroedinger anyons are considered. It is shown that for this special type of contact interactions scattering does not arise. By formulating the problem for the two particles system with a finite region of interaction which is then allowed to go to zero, it is established that this type of the above interactions is analog of the anomalous magnetic moment for fermions in the Aharonov-Bohm field. Finally, we consider connections with the Dirac anyon physics and Chern-Simons soliton solutions.

cond-mat/9501015

Electronic address: voropaev@tph100.physik.uni-leipzig.de

We suppose that the main ideas of the anyons as quasi-particles with fractional spin in (2+1) space-time are known to the readers. In other case, let us recommend [1] and [2] as general introduction with exhaustive bibliography. Contact interactions of anyons as point particles are considered in the works [3] and [4] in particular.

Really, we need for our consideration two facts only. Firstly, it is possible to describe anyon as a particle in an Aharonov-Bohm (AB) gauge field, which leads to the phase characterizing fractional statistics. Namely, we can analyze hamiltonian and wave function ($h = c = 1$) of the "free" point particle [3] ($V(\rho) = 0, \rho > 0$) either as

$$H = -\frac{1}{2M} \left[\frac{\partial^2}{\partial \rho^2} + \frac{1}{\rho} \frac{\partial}{\partial \rho} + \frac{1}{\rho^2} \frac{\partial^2}{\partial \phi^2} \right]; \Psi_\theta(\rho, \phi + 2\pi) = e^{i\theta} \Psi_\theta(\rho, \phi) \quad (1)$$

or as

$$H_\theta = -\frac{1}{2M} \left[\frac{\partial^2}{\partial \rho^2} + \frac{1}{\rho} \frac{\partial}{\partial \rho} + \frac{1}{\rho^2} \left(\frac{\partial}{\partial \phi} - i \frac{\theta}{2\pi} \right)^2 \right]; \Psi(\rho, \phi + 2\pi) = \Psi(\rho, \phi) \quad (2)$$

by means of a singular gauge transformation. Since the second hamiltonian is a separable operator, we can decompose the wave function $\Psi(\vec{\rho})$ as usual

$$\Psi(\vec{\rho}) = \sum_{m=-\infty}^{+\infty} C_m e^{im\phi} R_m(\rho) \quad (3)$$

so that the Schrödinger equation becomes

$$-\frac{1}{2M} \left[\frac{d^2}{d\rho^2} + \frac{1}{\rho} \frac{d}{d\rho} - \frac{(\nu_m)^2}{\rho^2} \right] R_m(\rho) = E R_m(\rho), \rho > 0 \quad (4)$$

with

$$\nu_m = \left| m - \frac{\theta}{2\pi} \right|; \nu = \theta/2\pi; 0 \leq \nu < 1 \quad (5)$$

For two identical anyons (ρ, ϕ) are the relative coordinates (in c.c.m.) and a rotation of π already leads to a physically indistinguishable state. So, we conclude that

$$\Psi_\theta(\rho, \phi + \pi) = e^{i\theta} \Psi_\theta(\rho, \phi) \quad (6)$$

and for radial wave function

$$\nu_m = \left| 2m - \frac{\theta}{\pi} \right|; \nu = \theta/\pi; 0 \leq \nu < 1 \quad (7)$$

analogous to the above consideration.

Secondly, in 2+1 - dimensions the equation for a "free" point Dirac anyon [4] is

$$(i\hat{\partial} - e\hat{A} - M)\Psi = 0 \quad (8)$$

where $\gamma^0 = \sigma^1, \gamma^1 = i\sigma^2, \gamma^2 = -i\sigma^1$ with an Aharonov-Bohm type magnetic field

$$A^i = -\frac{F}{2\pi} \varepsilon^{ij} \frac{\rho^j}{\rho}; B = \varepsilon^{ij} \partial_i A^j = F \delta(\vec{\rho}) \quad (9)$$

The iteration of the Dirac equation provides

$$[(-i\vec{\nabla} - e\vec{A})^2 - eB]\Psi^1 = (E^2 - M^2)\Psi^1; [(-i\vec{\nabla} - e\vec{A})^2 + eB]\Psi^2 = (E^2 - M^2)\Psi^2 \quad (10)$$

These components are very similar to the above wave function with the natural relativistic generalization : $E^2 \rightarrow E^2 - M^2$. But, this case of the minimal interaction is only a particular variant $\theta = \pi/2$ of all the possible self-adjoint extension of Dirac hamiltonian [5] general parametrization ($0 \leq \theta \leq \pi; eF > 0$). For arbitrary θ , contact interacting Dirac anyons satisfy coupled Schrödinger equations in the interaction region. Finally, let us note that the correct consideration of the above problem requires a regulated contact interaction or extended quasi-particles [4].

The main aim of this paper is the analysis of the possible contact interactions which provide a physical interesting behavior of anyons. One of them is the absence of self-scattering for one anyon and non-scattering on each other for two anyons, i.e. "soliton"-type behavior. Let us suppose that the interaction region is finite with radius $R = \text{Const}$ in spite of the above consideration. We investigate here non-relativistic particles for simplicity, but generalization for Dirac anyons is direct because of the remark that the components Ψ^1, Ψ^2 are decoupled outside the region of interaction.

So, the possible wave function of anyon (5),(6) is

$$\Psi^{(>)}(\vec{\rho}) = A \sum_{m=-\infty}^{+\infty} e^{im\Delta\phi} (-1)^m \left[e^{-i\pi\nu_m/2} J_{\nu_m}(p\rho) + e^{+i\pi\nu_m/2} J_{-\nu_m}(p\rho) \right], \rho > R \quad (11)$$

where

$$\Delta\phi \equiv \phi - \phi_\perp; (\vec{p}\vec{\rho}) = p\rho \cos(\phi - \phi_\perp) \quad (12)$$

It is easy to show by means of the integral representation [6]

$$\sum_{m=1}^{\infty} e^{im\theta} J_{m+\nu}(z) = \frac{1}{2} e^{iz \sin(\theta)} \int_0^z dx e^{-ix \sin(\theta)} \left[e^{i\theta} J_\nu(x) + J_{\nu+1}(x) \right] \quad (13)$$

that $\Psi^{(>)}(\vec{\rho})$ is

$$\Psi^{(>)}(\vec{\rho}) = A \sin(\pi\nu) \frac{1}{2} e^{i(\vec{p}\vec{\rho})} \int_0^\infty dx e^{-ix \cos(\Delta\phi)} \left[e^{+i\pi(1-\nu)/2} H_{1-\nu}^{(1)}(x) - e^{+i\Delta\phi} e^{+i\pi\nu/2} H_\nu^{(1)}(x) \right] \quad (14)$$

Using the following representation [7]

$$\int_0^\infty dx e^{-ixy} e^{+i\pi\nu/2} H_\nu^{(1)}(x) = \frac{2}{(1-y^2)^{1/2} \sin(\pi\nu)} \sin[\nu(\pi/2 + \arcsin y)], | \nu | < 1, | y | < 1 \quad (15)$$

we obtain the wave function of anyon

$$\Psi^{(>)}(\vec{\rho}) = -A e^{i\nu(\pi-\phi_\perp)} e^{i\nu\phi} e^{+i(\vec{p}\vec{\rho})} \frac{\sin(\Delta\phi)}{|\sin(\Delta\phi)|} \quad (16)$$

for the angles region $\Delta\phi \neq 0, \pi$ ($|\cos(\Delta\phi)| < 1$) and $0 \leq \Delta\phi < 2\pi$.

The analysis of the wave function for angles $\Delta\phi = 0, \pi$ is a more complex one. We mention shortly that expressions of the type

$$\int_0^\infty J_\nu(x) \begin{bmatrix} \cos x \\ \sin x \end{bmatrix} dx = (\infty \text{ or } 0) \quad (17)$$

are ill defined [7]. It prompts the idea that the wave function in this angle region is a generalized function analogous to the Dirac's δ -function. So, it is necessary to introduce some "classical" core for anyon. Finally, let us note that for the particular value $\nu = 1/2$ ("semion") and $\Delta\phi = 0$: $\Psi^{(>)}(\vec{\rho}) = 0$, i.e. complete reflection occurs at the origin.

Analogous, the wave function of the two-anyons system (in c.c.m.) (9) may be represented as

$$\Psi^{(>)}(\vec{\rho}) = A \sum_{m=1}^{\infty} e^{+i2m\Delta\phi} (-1)^m [J_{\nu_m}(p\rho) + BJ_{-\nu_m}(p\rho)] + \quad (18)$$

$$+ A \sum_{m=0}^{\infty} e^{-i2m\Delta\phi} (-1)^m [BJ_{\nu'_m}(p\rho) + J_{-\nu'_m}(p\rho)]; \quad (19)$$

where $\rho > R$, $\nu'_m = 2m + \theta/\pi$, $A(B) = \text{Const.}$ This expression provides quite natural wave functions for the free two-particles system in (a) the bosonic limit ($\nu = \theta/\pi = 0$):

$$\Psi_b^{(>)}(\vec{\rho}) = \left[\frac{1}{2} A(1+B) \right] [e^{+i(\vec{p}\vec{\rho})} + e^{-i(\vec{p}\vec{\rho})}]; \quad (20)$$

and (b) the fermionic limit ($\nu = \theta/\pi = 1$):

$$\Psi_f^{(>)}(\vec{\rho}) = e^{+i\nu\phi} \left[\frac{1}{2i} A(B-1) e^{-i\nu\phi_\perp} \right] [e^{+i(\vec{p}\vec{\rho})} - e^{-i(\vec{p}\vec{\rho})}] \quad (21)$$

On the other hand, $\Psi^{(>)}(\vec{\rho})$ (18) may be transformed by means of the integral representation [7]

$$\sum_{k=0}^{\infty} (t)^{2k} J_{2k+\nu}(z) = (z)^{-\nu} \int_0^z dx x^\nu J_{\nu-1}(x) \cosh\left(\frac{tx^2}{2z} - \frac{tz}{2}\right) \quad (22)$$

to the more appropriate form

$$\Psi^{(>)}(\vec{\rho}) = e^{+i\nu\phi} \left[\frac{1}{2} e^{-i\nu\phi_\perp} A \right] ([e^{-i\pi\nu/2} + Be^{+i\pi\nu/2}] e^{-i(\vec{p}\vec{\rho})} + \quad (23)$$

$$+ [e^{+i\pi\nu/2} + Be^{-i\pi\nu/2}] e^{+i(\vec{p}\vec{\rho})}); 0 < \phi < \pi; \rho > R \quad (24)$$

which coincides with the above boson and fermion wave functions for the corresponding limit cases.

Using the well-known "hard-core" requirement for the anyon interaction [2], we suppose that the "back-scattering" is equal to the "forward-scattering". So, we can conclude from the expression (23) that the wave function of the anyon before "scattering" is

$$\Psi^{(>)}(\vec{\rho}) = Ce^{-i[(\vec{p}\vec{\rho}) + (Et + \Delta/2)]} \quad (25)$$

and after "scattering" it will be

$$\Psi^{(>)' }(\vec{\rho}) = C e^{+i[(\vec{p}\vec{\rho}) - (Et - \Delta/2)]} \quad (26)$$

Here we used the condition $B \in \mathbf{R}$ for the "soliton"-type contact interaction. (Let us note that this condition provides self- adjointness of the initial hamiltonian also). It is easy to show that the phase shift Δ which appears after interaction of anyons with each other is determined by ν and B

$$tg(\Delta/2) = tg(\pi\nu/2) \frac{1-B}{1+B} \quad (27)$$

The physical interpretation of this result may be realised as a combined contact interaction of the two anyons which includes a statistic term, i.e. interaction of the Chern-Simons (CS) charge-flux, and term from the CS spin-flux interaction. So, using the general expression for the phase shift Δ (27) we can see that the boson (fermion) limit is possible now as when $\nu \rightarrow 0(1)$ as when $B \rightarrow 1(-1)$.

The remaining interesting problem for applications is the determination of the interaction potential between two "free" anyons. It require methods of the inverse scattering in general. We submit here a more simple approach which is valid when $(pR) \rightarrow 0$ only, i.e in the point quasi-particles limit. Then, we can approximate the above interaction potential by means of the Dirac's δ - function, namely $V_{int} \sim \lambda \delta(\rho/R - 1)$. Using the well-known boundary condition following from the properties of δ - function

$$R_m(1+0) = R_m(1-0); \frac{dR_m}{dx}(1+0) - \frac{dR_m}{dx}(1-0) = \lambda R_m(1); x = \rho/R \quad (28)$$

it is easy to show from (18) that for the s-wave ($m = 0$)

$$\lambda = -\nu + (pR/2)^{2\nu} \frac{2}{B} \frac{\Gamma(1-\nu)}{\Gamma(\nu)} \quad (29)$$

In continuation of the analogy with the particle in the AB field, we can add to the initial hamiltonian (1) or (4) the following term $V_{int} = -\mu_0 \frac{g}{2} \frac{F}{2\pi} \frac{1}{R} \delta(\rho - R)$ which describes the particle with anomalous magnetic moment (AMM) $a_e = \frac{g-2}{2}$ [8]. So, when $\mu_0 = e/2M; eF/2\pi = \nu$ we have for the attractive potential $\lambda = -[g/2]\nu = -[(g-2)/2 + 1]\nu$ and AMM will be determined by B

$$a_e = \frac{g-2}{2} = -(pR/2)^{2\nu} \frac{2}{B} \frac{\Gamma(1-\nu)}{\Gamma(1+\nu)} \quad (30)$$

For fermion with AMM in the δ - shell AB magnetic field $B = -1$ which coincides with the main result of [8].

It is clear now that we have considered the non-relativistic limit of the Dirac equation with the non-minimal coupling by means of the Pauli term type. On the other hand, in the relativistic Abelian Chern-Simons theory with the Lagrange density

$$L = \frac{k}{4} \epsilon^{\alpha\beta\gamma} A_\alpha F_{\beta\gamma} + (D_\mu \phi)^* (D_\mu \phi) - \frac{1}{k^2} | \phi |^2 (| \phi |^2 - v^2)^2 \quad (31)$$

topological solitons/vortices with quantized flux and non-topological solitons with no quantized flux are possible [9]

$$\nabla^2 \ln |\phi|^2 = -\frac{4v^2}{k^2} |\phi|^2 (1 - |\phi|^2/v^2) \quad (32)$$

The later are especially interesting for us because of the appropriate matter field distribution as in our δ -shell model

$$|\phi|^2 \rightarrow \rho^{2N-2}, \rho \rightarrow 0; |\phi|^2 \rightarrow \rho^{-2(\alpha+1)}, \rho \rightarrow \infty, \quad (33)$$

where $\alpha \geq N$, $N = 1, 2, \dots$ is a soliton number. These solutions possess angular momentum $J = kF/2(\alpha - N)$ where F is CS flux. One may consider the motion of the non-topological solitons and derive an effective Lagrangian from the underlying field theory. It is reasonable that as in topological soliton considerations [10] one finds a statistical interaction term with the value of the statistical factor in agreement with the generalized spin-statistic relation. So, it is possible that our consideration of the contact interaction of "free" anyons will provide a quantum version of the non-topological Abelian CS solitons.

I would like to thank Dr. M. Bordag, Dr. M. Hellmund for helpful conversations and the Institute Theoretical Physics, Leipzig University for kind hospitality. Also, I want to express my deep gratitude Dr. R. Herrmann for the support and as an initiator of a number of ideas.

References

- [1] F.Wilczek. "Fractional Statistics and Anyon Superconductivity" (World Scientific, Singapore, 1989).
- [2] A.Lerda. "Anyons" (Springer-Verlag, 1992).
- [3] C.Manuela, R.Tarrach. Phys.Lett.B, 222, v.268 (1991).
- [4] C.Manuela, R.Tarrach. Phys.Lett.B, 72, v.301 (1993).
- [5] Ph.de Sousa Gerbert. Phys.Rev.D., v.40, 1346 (1989).
- [6] V.Skarginsky. Trudi Lebedev Fiz.Inst. (Moscow), v.167, 142 (1987).
- [7] A.P.Prudnikov, Yu.A.Brychkov, and O.I.Marichev. "Integrals and Series, vol.2: Special Functions" (Gordon and Breach, New York, 1988).
- [8] M.Bordag, S.Voropaev. J.Phys.A., v.26, 7637 (1993); Phys. Lett. v.B333, 238 (1994).
- [9] R.Jackiw, S.-Y.Pi. Phys.Rev.D., v.42, 3500 (1990); CTP 2000, BU-HEP 91-19.
- [10] S.Kim, H.Min. Seoul National University preprint SNU TP 91-41 (1991).